

National  
Library  
of MedicineMy NC  
[Sign In] [Regis]

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Book

Search PubMed for [ ] Go Clear

Limits Preview/Index History Clipboard Details

Display Abstract Show 20 Sort by Send to

All: 1 Review: 0

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

My NCBI (Cubby)

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

1: Vet Immunol Immunopathol. 1987 Dec;17(1-4):303-12. Related Articles, Links

### Progress on gene transfer in farm animals.

Pursel VG, Rexroad CE Jr, Bolt DJ, Miller KF, Wall RJ, Hammer RE, Pinkert CA, Palmiter RD, Brinster RL.

Laboratory of Reproductive Physiology, University of Pennsylvania.

Transgenic pigs and sheep have been produced by the microinjection of single-cell zygotes and two-cell ova with linear molecules of mouse metallothionein I (MT) promoter/regulator fused to either the human growth hormone (hGH) or bovine growth hormone (bGH) structural genes. The foreign genes integrated into the chromosomes of 3 of 111 lambs or fetuses and 31 of 341 pigs or fetuses examined. Immunoreactive hGH or bGH was present in the plasma of two transgenic lambs and 19 transgenic pigs. The hGH concentration in plasma varied greatly among pigs and was unrelated to the number of gene copies that had integrated. Rate of growth was not enhanced in any of the transgenic pigs in comparison to their littermate controls. However, bGH and hGH exerted definite biological effects in transgenic pigs as evidenced by significantly depressed backfat measurements, elevated levels of insulin-like growth factor (IGF-I), stimulation of mammary development (by hGH) and reduction in porcine growth hormone (pGH) to nondetectable levels in plasma. Five of six founder transgenic pigs transmitted the MT-hGH gene construct to one or more progeny. Three progeny of a boar that expressed hGH also expressed the foreign gene.

PMID: 3481144 [PubMed - indexed for MEDLINE]

Display Abstract Show 20 Sort by Send to

[Write to the Help Desk](#)[NCBI | NLM | NIH](#)[Department of Health & Human Services](#)[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)